

Biocorrosion of materials and sick building syndrome

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 CSIRO. All rights reserved. The problem of biological damage of mineral building materials and structures based on them is multifaceted and covers all types of industry. The most destructive biocorrosion impacts are on building materials in cities with a large water area. Various types of microorganisms, including pathogens, and especially the filamentous fungi of the genera *Aspergillus*, *Penicillium*, *Trichoderma*, etc., occupy the surfaces of mineral building materials, cause their destruction, disturb the ecological balance of cities and endanger the human health. The term 'sick building syndrome' (SBS) is used to describe a situation when the residents of a building experience acute health- or comfort-related effects that seem to be linked directly to the time spent in the building wherein no specific illness or cause can be identified. Biological contaminants, in particular micromycetes, can present one of the possible causes of SBS. Here, we assessed the biodeterioration level of structural materials on the basis of fine-grained concrete widely used in construction practice and architecture. First, we determined the strength characteristics of the material that has been biologically damaged; second, we identified the damaging micromycetes and analysed their metabolic activity related both to the induction of biocorrosion and to the impacts of fungi on human health. Applying a new integrated approach, which combines methods of molecular microbiology and genetic toxicology with standard methods for determining the strength of building structures, we confirmed the relation between biodestructive and pathogenic properties of micromycete isolates.

<http://dx.doi.org/10.1071/MA18040>

References

- [1] Joshi, S. M. (2008) The sick building syndrome. *Indian J. Occup. Environ. Med.* 12, 61-64. doi:10. 4103/0019-5278. 43262
- [2] Haleem Khan, A. A. And Mohan Karuppayil, S. (2012) Fungal pollution of indoor environments and its management. *Saudi J. Biol. Sci.* 19, 405-426. doi:10. 1016/ j. sjbs. 2012. 06. 002
- [3] Gomzi, M. And Bobic, J. (2009) Sick building syndrome. Do we live and work in unhealthy environment *Periodicum Biologorum* 111, 79-84.
- [4] van Burik, J.-A. H. And Magee, P. T. (2001) Aspects of fungal pathogenesis in humans. *Annu. Rev. Microbiol.* 55, 743-772. doi:10. 1146/annurev. micro. 55. 1. 743
- [5] Allsopp, D. et al. (2004) *Introduction to biodeterioration*. 2nd edn. Cambridge University Press.
- [6] Jayakumar, S. (2012) *Studies on the biodeterioration of concrete by marine algae*. Shodhganga Repository of Indian Electronic Theses and Dissertations. Pondicherry University School of Engineering. [http://shodhganga.inflibnet. Ac. in/handle/ 10603/5590](http://shodhganga.inflibnet.ac.in/handle/10603/5590)
- [7] Wei, S. et al. (2013) Microbiologically induced deterioration of concrete. *Braz. J. Microbiol.* 44, 1001-1007. doi:10. 1590/S1517-83822014005000006
- [8] Bertron, A. et al. (2004) Cement paste alteration by liquid manure organic acids: chemical and mineralogical characterization. *Cement Concr. Res.* 34, 1823-1835. doi:10. 1016/j. cemconres. 2004. 01. 002

- [9] Park, M. et al. (2013) Lipolytic enzymes involved in the virulence of human pathogenic fungi. *Mycobiology* 41, 67-72. doi:10. 5941/MYCO. 2013. 41. 2. 67
- [10] Yike, I. (2011) Fungal proteases and their pathophysiological effects. *Mycopathologia* 171, 299-323. doi:10. 1007/s11046-010-9386-2
- [11] Vylkova, S. (2017) Environmental pH modulation by pathogenic fungi as a strategy to conquer the host. *PLoS Pathog.* 13, e1006149. doi:10. 1371/journal. ppat. 1006149
- [12] Yakovleva, G. et al. (2016) Assessment of biodamage resistance of various concrete grades. *Int. J. Pharm. Technol.* 8, 24291-24299.
- [13] Aleksic, B. et al. (2017) Aerosolization of mycotoxins after growth of toxinogenic fungi on wallpaper. *Appl. Environ. Microbiol.* doi:10. 1128/AEM. 01001-17
- [14] Heutte, N. et al. (2017) Assessment of multi-contaminant exposure in a cancer treatment center: A 2-year monitoring of molds, mycotoxins, endotoxins, and glucans in bioaerosols. *Environ. Monit. Assess.* 189, 31. doi:10. 1007/s10661-016-5751-z
- [15] Zhang, J. et al. (2016) A polyketide synthase encoded by the gene An15g07920 is involved in the biosynthesis of ochratoxin A in *Aspergillus niger*. *J. Agric. Food Chem.* 64, 9680-9688. doi:10. 1021/acs. jafc. 6b03907